

AMENDMENT TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

1-14. (canceled)

15. (Currently amended) A method for the inhibition of transient lower esophageal sphincter relaxations (TLESRs), **the method comprising administering** whereby a **therapeutically** pharmaceutically and pharmacologically effective amount of a **compound which is a** metabotropic glutamate receptor 5 antagonist, or a pharmaceutically acceptable salt **of the compound**, or an optical isomer **of the compound or a pharmaceutically acceptable salt of the optical isomer**, thereof, is administered to a **patient suffering from gastroesophageal reflux disease** subject in need of such inhibition.

16. (Currently amended) A method for the treatment of gastro-esophageal reflux disease (GERD), **the method comprising administering** whereby a **therapeutically** pharmaceutically and pharmacologically effective amount of a **compound which is a** metabotropic glutamate receptor 5 antagonist, or a pharmaceutically acceptable salt **of the compound**, or an optical isomer **of the compound or a pharmaceutically acceptable salt of the optical isomer**, thereof, is administered to a **patient suffering from gastroesophageal reflux disease** subject in need of such treatment.

17. (Currently amended) A method for the **inhibition** prevention of reflux **of gastric juice**, **the method comprising administering** whereby a **therapeutically** pharmaceutically and pharmacologically effective amount of a **compound which is a** metabotropic glutamate receptor 5 antagonist, or a pharmaceutically acceptable salt **of the compound**, or an optical isomer **of the compound or a pharmaceutically acceptable salt of the optical isomer**, thereof, is administered to a **patient suffering from gastroesophageal reflux disease** subject in need of such prevention.

18. (Currently amended) A method for the treatment of, ~~or prevention of~~, regurgitation of gastric juice, the method comprising administering whereby a therapeutically pharmaceutically and pharmacologically effective amount of a compound which is a metabotropic glutamate receptor 5 antagonist, ~~or a pharmaceutically acceptable salt of the compound,~~ or an optical isomer of the compound or a pharmaceutically acceptable salt of the optical isomer, thereof, is administered to a patient suffering from gastroesophageal reflux disease ~~subject in need of such inhibition.~~

19. (Withdrawn) A method for the prevention of, or treatment of, lung disease, whereby a pharmaceutically and pharmacologically effective amount of a metabotropic glutamate receptor 5 antagonist, or a pharmaceutically acceptable salt or an optical isomer thereof, is administered to a subject in need of such treatment or prevention.

20. (Withdrawn) A method for managing failure to thrive, whereby a pharmaceutically and pharmacologically effective amount of a metabotropic glutamate receptor 5 antagonist, or a pharmaceutically acceptable salt or an optical isomer thereof, is administered to a subject in need of such management.

21. (Withdrawn) A method for treatment or prevention of asthma, whereby a pharmaceutically and pharmacologically effective amount of a metabotropic glutamate receptor 5 antagonist, or a pharmaceutically acceptable salt or an optical isomer thereof, is administered to a subject in need of such treatment or prevention.

22. (Withdrawn) A method according to claim 21, wherein the asthma is reflux-related asthma.

23. (Withdrawn) A method for treatment or prevention of chronic laryngitis, whereby a pharmaceutically and pharmacologically effective amount of a metabotropic glutamate receptor 5 antagonist, or a pharmaceutically acceptable salt or an optical isomer thereof, is administered to a subject in need of such treatment or prevention.

24. (Currently amended) **The [A]** method according to any one of claims 15-23, wherein the metabotropic glutamate receptor 5 antagonist is 2-methyl-6-(phenylethynyl)-pyridine.

25. (Currently amended) **The [A]** method according to claim 24, wherein the metabotropic glutamate receptor 5 antagonist is the hydrochloride salt of 2-methyl-6-(phenylethynyl)-pyridine.

26. (Withdrawn) A method according to any one of claims 15-23, wherein the metabotropic glutamate receptor 5 antagonist is 3-[3-(5-fluoropyridin-2-yl)-1,2,4-oxadiazol-5-yl]-5-(methoxymethyl)benzonitrile.

27. (Withdrawn) A method according to any one of claims 15-23, wherein the metabotropic glutamate receptor 5 antagonist is 3-fluoro-5-[3-(5-fluoropyridin-2-yl)-1,2,4-oxadiazol-5-yl]benzonitrile.

28. (Currently amended) **The [A]** method according to any one of claims **15-23** [15-27], wherein the daily dose of the metabotropic glutamate receptor 5 antagonist is from 0.1-100 mg per kg body weight of the subject to be treated.